



## Addendum Addendum to “Groups of ribbon knots”

Ka Yi Ng\*

145 Lincoln Avenue, Apt. 5W, Staten Island, NY 10306, USA

Received 2 October 2003; accepted 19 July 2004

---

### Abstract

The inductive step that was described in the proof of Theorem 3.2 of Ng (Topology 37(2) (1998) 441) is clarified.  
© 2004 Published by Elsevier Ltd.

*Keywords:* Vassiliev invariants; Ribbon knots

---

The purpose of this document is to clarify the inductive step described in the proof of Theorem 3.2 in [2]. In the second last sentence of the proof, it says, ‘it follows from the inductive proof that  $\mathcal{R}_n$  is of index two.’ The question of how this assertion is verified was first raised by Natan and his student Ron [1]. To avoid any confusions that may arise in the future, the author of the paper [2] would like to fill in the details to show that  $\mathcal{R}_n$  is of index two. Readers of this document are assumed to have familiarity with [2]. For your reference, Theorem 3.2 is stated below.

**Theorem 3.2.**  *$\mathcal{R}_n$  forms a subgroup of the free abelian group  $\mathcal{V}_n$  of index two. So its rank is the same as the rank of  $\mathcal{V}_n$  and is the number of linearly independent primitive rational invariants of order  $\leq n$ .*

To show that  $\mathcal{R}_n$  is of index two, we shall establish the following two lemmas.

**Lemma A.** *Let  $D_1, \dots, D_r$  be all the distinct non-split chord diagrams of order  $n \geq 2$ . Let  $\lambda_1, \dots, \lambda_r$  be  $r$  arbitrary integers. Then for  $n > 2$ , there exists a ribbon knot  $K_n$  whose additive invariants of order  $< n$*

---

DOI of original article 10.1016/S0040-9383(97)00037-2.

\* Tel.: +1-718-987-9351.

E-mail addresses: [kaying7@hotmail.com](mailto:kaying7@hotmail.com), [kayi.ng@wallstreetsystems.com](mailto:kayi.ng@wallstreetsystems.com) (K.Y. Ng).

are trivial and  $V(K_n) = \sum_{i=1}^r \lambda_i V(D_i)$  for all additive invariants  $V$  of order  $n$ . When  $n = 2$ , there exists a ribbon knot  $K_2$  with  $V(K_2) = 2\lambda_1 V(D_1)$  for all additive invariants  $V$  of order 2.

**Lemma B.** *Given a knot  $K$  whose additive invariants of order  $< n$  are trivial, there exist  $n$ -chord diagrams  $D_1, \dots, D_r$  and integers  $\lambda_1, \dots, \lambda_r$  such that  $V(K) = \sum_{i=1}^r \lambda_i V(D_i)$  for all additive invariants  $V$  of order  $n$ .*

Lemma A is an extended version of Lemma 3.1 in [2] and can be proved using the same arguments as given in [2]. Lemma B is proved in [3]. We shall now make use of these two lemmas to prove Lemma C.

**Lemma C.** *Let  $X$  be the right-handed trefoil. Let  $K$  be an arbitrary knot and let  $n \geq 2$  be an integer. Define  $s$  to be 1 if the Arf invariant of the knot  $K$  is non-zero and 0 otherwise. Then one can construct a ribbon knot  $K_n$  so that  $V(K \# sX \# K_n) = 0$  for all additive invariants  $V$  of order  $\leq n$ .*

**Proof of Lemma C.** For  $n = 2$ , the knot can always be represented by an integral multiple  $\lambda$  of the non-split 2-chord diagram  $D$  so that  $V(K) = \lambda V(D)$  for all additive invariants of order 2. Write  $\lambda$  as  $2p - s$  where  $p$  is an integer. By Lemma A, we can find a ribbon knot  $K_2$  with  $V(K_2) = -2pV(D)$  for all additive invariants of order 2. The right-handed trefoil knot has  $V(X) = V(D)$  and thus, the case  $n = 2$  is proved.

Suppose a ribbon knot  $K_n$  is constructed so that  $V(K \# sX \# K_n) = 0$  for all additive invariants of order  $\leq n$ . Lemma B tells us that there exist integers  $\lambda_1, \dots, \lambda_r$  and  $(n+1)$ -chord diagrams  $D_1, \dots, D_r$  so that  $V(K \# sX \# K_n) = \sum_{i=1}^r \lambda_i V(D_i)$  for all additive invariants of order  $n+1$ . Apply Lemma A to construct a ribbon knot  $R$  whose additive invariants of order  $\leq n$  are trivial and  $V(R) = -\sum_{i=1}^r \lambda_i V(D_i)$  for all additive invariants of order  $n+1$ . Then  $V(K \# sX \# K_n \# R) = 0$  for all additive invariants of order  $\leq n+1$ .

Lemma C implies that  $V(K \# sX \# K_n) = 0$  for all rational-valued invariants of order  $\leq n$ . According to the proof of Theorem 3.2 in [2],  $\mathcal{R}_n$  is a subgroup of  $\mathcal{V}_n$ . Thus, as an element in  $\mathcal{V}_n$ , we have  $[K]_n \in -[sX]_n + \mathcal{R}_n$ . This completes the proof that  $\mathcal{R}_n$  is of index two.  $\square$

The author would like to thank Joan Birman for help in the preparation of this note. Many thanks also go to Dror Bar Natan and his student Ofer Ron for raising the question and for numerous discussions.

## References

- [1] D.B. Natan, O. Ron, A correction to “Groups of Ribbon Knots” by KaYi Ng, arXiv:math.GT/0309282, 17 September 2003.
- [2] K.Y. Ng, Groups of ribbon knots, Topology 37 (2) (1998) 441–458.
- [3] K.Y. Ng, T. Stanford, On Gusarovs groups of knots, Proc. Cambridge Philos. Soc. 126 (1) (1999) 63–76.